ARTÍCULO ORIGINAL - ORIGINAL ARTICLE

Study of *Araucaria angustifolia* nuts by Gamma Spectrometry and X-Ray Fluorescence

Estudio de semillas de Araucaria angustifolia por Espectrometría Gamma y Fluorescencia de Rayos X

Edher Zacarías Herrera¹, Avacir Casanova Andrello¹, Carlos Roberto Appoloni¹

State University of Londrina, Department of Physics, Londrina, Paraná, Brazil.

Autor de correspondencia: edher712@gmail.com, acandrello@uel.br

DOI: https://doi.org/10.32480/rscp.2019-24-2.262-273

Recibido: 25/09/2019. Aceptado: 2/12/2019.

Abstract: The seed of Araucaria angustifolia is largely produced and consumed in Parana state, Brazil. It has an estimated annual consumption of approximately 0.167 kg per capita. The samples of seeds were acquired in 8 different commercial points of Londrina city in the state of Paraná, Brazil. To determine the natural level of radiation and elemental composition were analysed the seed pulp and peels, to determine the level of radiation in these samples and to calculate the effective dose per intake, the gamma ray spectrometry was applied using the HPGe detector with a 60% relative efficiency, and the X-rays Fluorescent by Dispersive Energy (EDXRF) was applied for elemental identification of the composition with a spectrometer from Shimadzu EDX-720 model. The samples were qualitatively analyzed by EDXRF, where the elements K, Fe, Cu, Zn, Rb, Ca, P, Mg and Sr were found in pulp and peels of seed, however, the elements Mg and Sr were detected only in the peel samples. The results obtained by gamma spectrometry showed that the radioactive activities in the Araucaria angustifolia pulp per kilogram in nature, for 226Ra ranged from 0.691 to 2.269 Bg/kg, the interval activity measured for the 228Ra were 0.386 to 1.767 Bg/kg and the 40K were 283.405 to 1,536.820 Bq/kg. Activities of 137Cs were detected in the pulp and peels of seed. The average dose per intake was 0.68 µSv/year for 226Ra and 0.35 µSv/year for 228Ra.

Keywords: Araucaria angustifolia, EDXRF, dose, radiation.

Resumen: La semilla de Araucaria angustifolia es producida y consumida en gran medida en el estado de Paraná, Brasil. Tiene un consumo anual estimado aproximado de 0.167 kg per cápita. Las muestras de semillas fueron adquiridas en 8 puntos comerciales diferentes de la ciudad de Londrina en el



estado de Paraná, Brasil. Para determinar el nivel de radiación natural y la composición elemental, se analizaron la pulpa y las cáscaras de las semillas, para determinar el nivel de radiación y dosis efectiva por ingesta, se aplicó la espectrometría de rayos gamma utilizando el detector HPGe con un 60% eficiencia relativa, se utilizó fluorescencia de rayos X por energía dispersiva (EDXRF) para la identificación elemental de la composición con un espectrómetro fabricado por Shimadzu modelo EDX-720. Las muestras fueron analizadas cualitativamente por EDXRF, donde los elementos K, Fe, Cu, Zn, Rb, Ca, P, Mg y Sr se encontraron en la pulpa y cascara de las semillas, sin embargo, los elementos Mg v Sr se detectaron solo en las cascaras de la semilla. Los resultados obtenidos por espectrometría gamma muestran que las pulpas de Araucaria angustifolia in natura por kg, poseen actividades de 226Ra en el rango fue 0.69 a 2.26 Bg/kg, para el 228Ra el rango fue de 0.38 a 1.76 Bg/kg, para el 40K el rango de actividad fue de 283.40 a 1,536.82 Bg/kg. Se detectaron actividades de 137Cs en la pulpa y las cáscaras de semillas. La dosis promedio por ingesta fue de 0,68 μSv/año para el 226Ra y 0.35 μSv/año para el 228Ra.

Palabras clave: Araucaria angustifolia, EDXRF, dosis, radiación.

1. INTRODUCTION

Radioactivity is present in all moments of our daily life. The radiation that exists in nature without human intervention is called natural radioactivity and it is present in the ground, in the air, etc. The radioactivity generated by human intervention is called artificial radioactivity (1-5). Its applications have increased in the last few years. Therefore, it is necessary a periodical environmental control to monitor human activities such as the analysis of different types of food in order to guarantee their quality (6-13).

The pulp of the araucaria's seed is suitable for human consumption, it can be ingested naturally, roasted or boiled, the peel is used in medical infusions for the treatment of muscle tension of varicose vein. In addition, the araucaria's resin is used for the treatment of respiratory infections and the leaves to treat scrofula, fatigue, and anemia (14–19). The commercialized volume of Araucaria nuts reached over 10 thousand tons of Araucaria nuts from August 1994 to June 2002 by the Parana City Supply Centers (CEASA, in Portuguese) from Curitiba, Maringá, Londrina, Foz do Iguaçu and Cascavel, which corresponds to an approximated average of 1300 ton per year (20).

The research often focuses on the nutritional content, chemistry characterization and some physicochemical properties of the Araucaria nuts (6, 7). However, due to its wide distribution and high amount of consumption,

there is a concerning risk of exposure to natural radioactivity from the Araucaria nuts. This study aims to determine the activity of ²³²Th, ²³⁸U, ⁴⁰K, and ¹³⁷Cs and to perform a multi-element analysis of the seed of Parana pine tree (*Araucaria Angustifolia*), commonly known as Pinhão (Araucaria nut) (15, 21).

2. MATERIALS AND METHODS

2.1. Preparation of samples

The samples were acquired in the city of Londrina (Parana, Brazil) between May and July of 2014. Among 8 batches were bought in different commercial centers and informal stores found around the city. During the purchase of Araucaria nuts, their origin and time of harvest were consulted, however, some traders could not specify this information.

The Araucaria nuts were roasted on a hot plate between 100°C and 150°C, simulating cooking procedures for human consumption. Later, their pulp and peel were separated, grounded manually in a mortar and stored in neutral plastic containers of 300mL of volume. At the end, 16 containers were used for the seed's pulp and 8 containers for the seed's peel which were sealed for 40 days in order to achieve the secular balance for the upcoming analysis.

For the X-ray fluorescence analysis, the pulp was dehydrated, sieved and pressed into tablets to achieve homogeneous samples. A binding agent (0.250 g of boric acid - H3BO3) was used for every 1000 g of seed powder. Two 2.5 mm thick tablets were prepared for each batch by applying 9 tons of uniaxial pressure over the mix for 2 minutes. In addition, the powder samples were also measured for elementary identification of the composition, putting 1 gram of Araucaria pulp powder and the peels over a chemplex (samples holder).

2.2. Gamma-ray spectrometry

A high-resolution gamma-ray spectrometer system setup was a HPGe (hyperpure germanium) detector, manufactured by CANBERRA (model GC6020) with a relative efficiency of 60% and the shield elaborated by ORTEC model HPLBS1 designed for environmental applications. This setup was used to measure the radionuclides of interest.

All spectra were collected during a 24-hour period, preceded by the acquisition of the background spectrum produced by natural radiation for the calculation of the activity and committed effective doses.

2.2. Energy dispersive X-ray fluorescence spectrometry (EDXRF)

A commercial equipment of X-ray fluorescence by dispersive energy (EDXRF) manufactured by Shimadzu (model EDX-720) with an Rh target (X-rays source) was used for the qualitative elemental analysis of the Araucaria nuts. A solid-state detector was used with the following experimental set up parameters: 10 mm collimator, 100 s of measurement time, 50 kV of voltage and 487 – 604 μA of current.

EDXRF analysis was performed for the elemental identification of the composition of the araucaria seeds and control of contamination with heavy metals. Because of the lack of certified material, it was not possible to quantify elements.

2.3. Estimation of the consumption per capita

Information and analysis about Araucaria nuts consumption per capita was not found in the literature. Due to this fact, the value was estimated through the quantity of Araucaria nuts commercialized among the different supply centers from Parana State and the Brazilian Institute of Geography and Statistics (IGBE, in Portuguese) shown in Table 1.

Table 1: Commercialized Araucaria nuts among the different supply centers of Parana State (22).

| District or city | Quantity (ton) | Population | Consumption per capita |
|------------------|-------------------|------------|------------------------|
| Curitiba | 511.04 | 1751907 | 0.292 |
| Cascavel | 106.1 | 286205 | 0.371 |
| Foz do Iguaçu | 30.25 | 256088 | 0.118 |
| Londrina | 14.68 | 357077 | 0.041 |
| Maringá | 14.68 | 357077 | 0.012 |
| | | Average | 0.167 |

The official figures might contain considerable discrepancy because of the informal or clandestine sales in rural areas, and because not all the Araucaria nuts sold, is consumed (23).

3. RESULTS AND DISCUSSIONS

3.1. Quantitative results

Figure 1 presents gamma spectrum pulps of araucaria, obtained with a 24 hours measure time. Peaks of interest from radionuclides are indicated. Because it is the most abundant in nature, the peak belonging ⁴⁰K is the most intense in comparations with others peaks.




Figure 1: A) Gamma acquisition system scheme and typical spectrum of araucaria B) peels and C) pulp.

Using the spectra obtained in gamma spectrometry, the Bq / kg activities of 226 Ra, 228 Ra, 40 K, and 137 Cs in the araucaria seed and bark samples were calculated.

Due to the different ways of consumption of Araucaria seeds the activities are calculated per kg in nature.

The radioactivity of the desired radionuclides for each sample is presented in Table 2. This table organizes the average activity of ²²⁶Ra, ²²⁸Ra, ⁴⁰K, and ¹³⁷Cs from 16 pulp samples and 8 Araucaria nuts samples *in nature* per kilogram.

Table 2: The average activity of 226 Ra, 228 Ra, 40 K, and 137 Cs per kilogram for pulp and peel of Araucaria nuts *in nature*.

| Danidana d Canada | Average activity (Bq/kg) | | | |
|-------------------|--------------------------|-------------------|-----------------|-------------------|
| Powdered Samples | ²²⁶ Ra | ²²⁸ Ra | ⁴⁰ K | ¹³⁷ Cs |
| Pulp | 0.93 ± 0.05 | 0.62 ± 0.02 | 498 ± 1 | 0.65 ± 0.04 |
| Peel | 5.8 ± 0.4 | 3.9 ± 0.2 | 730 ± 6 | 2.4 ± 0.4 |

The radionuclides_analyzed presented higher levels of activity in the Araucaria nuts peel. In all 8 batches of pulp of Araucaria nuts purchased, the presence of ¹³⁷Cs was registered, but just in the batch number 6 was registered the presence of ¹³⁷Cs in the peel.

In Figure 2 it is observed that all batches shown an activity of 137 Cs and 228 Ra, but the activity of 226 Ra is similar in all samples. The 137 Cs is absorbed by the tree roots due to similar properties that own the K (24–26), the activities of 37 Cs in all samples are below the concentration level of 600 Bq/kg for the sum of activities of 134 Cs and 137 Cs respectively.




Figure 2. The activity of ²²⁶Ra, ²²⁸Ra, ⁴⁰K, and ¹³⁷Cs for each batch of pulp of Araucaria nuts per kilogram *in nature*.

Has been calculated the effective dosage committed by ingestion (DEC) with the average of 226 Ra and 228 Ra and average Commercialized Araucaria nuts

among the different supply centers of Parana of araucaria nuts. Using the following expression for the calculated (DEC) (27–29).

$$Dec=e(g)A.C$$
 [1]

Where C is the consumption per capita in Kg/years, A is the activity of radionuclide Bq/kg and e(g) Is the effective dose compromised per unit in Sv/Bq.

The Table 3 shows the effective dosage committed by ingestion (*Dec*) with the UNSEAR reference value.

Table 3: Effective dosage committed by ingestion (DEC) calculated with the average of ²²⁶Ra and ²²⁸Ra.

| | Committed effective doses (μSv/year) | | | | | |
|----------------|--------------------------------------|-------------------|------|------------------------|-------------------|--------|
| Age (years) | Araucaria consumption | | nuts | Reference [UNSCEAR] | | values |
| | ²²⁶ Ra | ²²⁸ Ra | Sum | ²²⁶ Ra | ²²⁸ Ra | Sum |
| 2 a 7 | 0.96 | 0.58 | 1.54 | 7.5 | 31 | 38.5 |
| 7 a 12 | 0.8 | 0.39 | 119 | 12.0 | 40.0 | 52.0 |
| >17 | 0.28 | 0.07 | 0.35 | 6.3 | 11 | 17.3 |
| Average | 0.68 | 0.35 | 1.03 | 8.1 | 21 | 29.1 |

According to the average values of effective doses per radionuclide, it was verified that the ^{228}Ra present on the effective committed dose of 0.35 $\mu Sv/year$, which is two orders of magnitude lower than the reference values of UNSCEAR, 21 $\mu Sv/year$. For ^{226}Ra , the average effective dose weighted was 0.68 $\mu Sv/year$, approximately 13 times lower than the one of reference of 8.1 $\mu Sv/year$. The differences obtained for the doses and the reference values are expected since the calculations in the current research refer only to the dose of the Araucaria nuts consumption, while the reference values contain other daily diet food.

Furthermore, the *DEC* for 40 K was not calculated since the dose of this radionuclide is controlled by the organism through homeostatic control. The annual equivalent doses for tissues from 40 K are 165 and 185 μ Sv/year for adults and children, respectively. The effective dose is 170 μ Sv (29, 30).

3.2. Qualitative results by EDXRF

Figure 3 shows typical spectra of X Ray Fluorescence spectra on the Araucaria pulp and shield powdered. In all the pulp powder samples was detected the presence of P, S, K, Ca, Fe, Rb, Zn and Cu. However, in the pulp tablets, the frequency of Zn and Cu detection was lower than the frequency in the pulp.




Figure 3: Araucaria shield and pulp powders X-ray fluorescence spectra.

In the Table 3 shows element intensity of major elements present in the araucaria measurements with X-ray fluorescence and standard deviation in the Araucaria nuts powdered tablet, pulp and peel. No significant difference between powder and tablet peaks intensity was observed, except for the increased presence of Zn in the powdered pulp.

Table 3: Major elements intensity detected in Araucaria nuts powdered pulp, tablet, and peel.

| | | Intensity | |
|---------|-------------------|--------------------|--------------------|
| Element | Peel | Pulp Powder Tablet | Pulp powder |
| K | 1.865±0.081 | 3.417±0.635 | 3529±0.626 |
| S | 0.039 ± 0.001 | 0.073±0.003 | 0.074 ± 0.003 |
| Fe | 0.879 ± 0.014 | 0.83 ± 0.054 | 0.43 ± 0.02 |
| Cu | 0.612±0.014 | 0.663±0.024 | 0.707±0.019 |
| P | 0.27 ± 0.011 | 0.051±0.005 | 0.0495 ± 0.005 |
| Ca | 0.34 ± 0.014 | - | - |
| Mn | 0.29±0.013 | - | - |
| Rb | - | 3.940±0.106 | 3.714±0.061 |
| Zn | - | - | 0.587 ± 0.014 |

In all the pulp powder samples was detected the presence of P, S, K, Ca, Fe, Rb, Zn and Cu. However, in the pulp tablets, the frequency of Zn and Cu detection was lower than the frequency in the pulp.

P, K, Rb, Fe, S, Ca, Cu and Zn were detected both in the pulp and the peels. Sr and Mn were observed only in the peels. No trace of Sr was detected in the pulp and the peel is consumed as a medicinal tea. As the Sr is not soluble in water and as the consumption is not frequent, this practice does not generate a risk.

The Cu is a transition non-ferrous metal abundant on earth, however, our interaction with this element is not daily. It is present in almost all types of food and in the air. Considered a micronutrient, Cu is only dangerous at high concentration and long periods of exposition, in which cases, it can cause damage to the liver and the kidney. It was not yet confirmed if Cu is a carcinogenic element (31, 32).

4. CONCLUSIONS

Araucaria nuts are not consumed in high amounts by the population, they are a typical food but not popular in Brazil.

The values of Dec calculated by gamma spectrometry show that the consumption of araucaria nuts are in acceptable values of 226 Ra and 228 Ra, in addition to verifying that the levels of activity of 137 Cs and 46 K are in the range accepted by UNSCEAR(33).

Rev. Soc. cient. Parag. 2019;24(2):262-273

ACKNOWLEDGMENTS

This work was supported by the *Brazilian* National Council for Scientific and Technological Development – CNPq.

REFERENCES

- 1. Knoll GF, Kraner HW. Radiation Detection and Measurement. Proceedings of the IEEE. 1981.
- 2. Desideri D, Meli MA, Roselli C. Natural and artificial radioactivity determination of some medicinal plants. J Environ Radioact. 2010;101(9):751-6.
- 3. Peirson DH. Artificial radioactivity in Cumbria: summary of an assessment by measurement and modelling. J Environ Radioact. 1988;6(1):61–75.
- 4. Steinhauser G, Merz S, Hainz D, Sterba JH. Artificial radioactivity in environmental media (air, rainwater, soil, vegetation) in Austria after the Fukushima nuclear accident. Environ Sci Pollut Res. 2013;20(4):2527–34.
- 5. Baggoura B, Noureddine A, Benkrid M. Level of natural and artificial radioactivity in Algeria. Appl Radiat Isot. 1998;49(7):867–73.
- 6. Antovic I, Antovic NM. Determination of concentration factors for Cs-137 and Ra-226 in the mullet species Chelon labrosus (Mugilidae) from the South Adriatic Sea. J Environ Radioact, 2011.
- 7. Shcheglov AI, Tsvetnova OB, Klyashtorin A. The fate of Cs-137 in forest soils of Russian Federation and Ukraine contaminated due to the Chernobyl accident. J Geochemical Explor. 2014;142:75–81.
- 8. Galeriu D, Oncescu M. Environmental radioactivity of 1987 in Romania. Central Inst. of Physics, 1988.
- 9. Karakelle B, Öztürk N, Köse A, Varinlioğbrevelu A, Erkol AY, Yilmaz F. Natural radioactivity in soil samples of Kocaeli basin, Turkey. J Radioanal Nucl Chem. 2002;254(3):649–51.
- 10. Xiao T, Zhao Z. The investigation of environmental radioactivity background around a pulsed reactor. Nucl Power Eng. 1990;11(3):77–9.
- 11. VandenBygaart AJ, Protz R. Gamma radioactivity in podzolic soils of Northern Ontario, Canada. J Environ Radioact. 1999;42(1):51–64.
- 12. Schauble EA, Ghosh P, Eiler JM. Preferential formation of 13C–180 bonds in carbonate minerals, estimated using first-principles lattice dynamics. Geochim Cosmochim Acta. 2006;70(10):2510–29.
- 13. Femandes EAN. Environmental monitoring in the uranium mining and milling facility at pocos de caldas. Nucl Phys Prot. 1995;86.
- 14. Rezende SC de. Valorização da casca do pinhão, um subproduto da semente de Araucaria angustifolia, para produção de materiais poliméricos. Vol. 1, The British Journal of Psychiatry. Instituto Politécnico de Bragança; 2016.

- 15. Tavares Bastos Gama TMM, Masson ML, Haracemiv SMC, Zanette F, Córdova KRV. A influência de tratamentos térmicos no teor de amido, colorimetria e microscopia de pinhão nativo (araucaria angustifólia) e pinhão proveniente de polinização controlada. Rev Bras Tecnol Agroindustrial [Internet]. 2010 Dec 17 [cited 2018 Dec 14];4(2). Available from: https://periodicos.utfpr.edu.br/rbta/article/view/592
- 16. Silva LACS, Modolo DM, Martinez P, Piero EA di, Bigide P, Arthur V, et al. Effects of gamma radiation and storage on cooked pine seed (Araucaria angustifollia), 2011;
- 17. Zandavalli RB, Dillenburg LR, de Souza PVD. Growth responses of Araucaria angustifolia (Araucariaceae) to inoculation with the mycorrhizal fungus Glomus clarum. Appl Soil Ecol. 2004;25(3):245–55.
- 18. Leite DM da C. Avaliação nutricional da semente do Pinheiro-do-Paraná (Araucaria angustifolia). 2007;
- 19. Oubari official Website [Internet]. 2015 [cited 2019 Sep 15]. Available from: https://www.ppmac.org/content/pinhão-araucária
- 20. Balbinot R, Garzel JCL, Weber KS, Ribeiro AB. Tendências de consumo e preço de comercialização do pinhão (semente da Araucaria angustifolia (Bert.) O. Ktze.), no estado do Paraná. Ambiência, 2008;
- 21. Gama TMMTB. Estudo comparativo dos aspectos físico-químicos do pinhao nativo e do pinhao proveniente de processos de polinização controlada de Araucaria angustifolia e da influencia do tratamento térmico, 2013.
- 22. Instituto Brasileiro de Geografia e Estatística. Censo I. Available from: http://www.censo2010. ibge.gov.br/
- 23. Balbinot R, Carlos J, Garzel L, Weber KS, Ribeiro AB. Trends of consumption and commercialization price of the Brazilian-pine nut-seed of Araucaria angustifolia (Bert.) O. Ktze., in the state of Parana [Internet]. [cited 2018 Dec 13]. Available from: https://revistas.unicentro.br/index.php/ambiencia/article/viewFile/147/165
- 24. Maathuis FJM, Sanders D. Mechanisms of potassium absorption by higher plant roots. Physiol Plant. 1996;96(1):158–68.
- 25. Johansen C, Edwards DG, Loneragan JF. Potassium fluxes during potassium absorption by intact barley plants of increasing potassium content. Plant Physiol. 1970;45(5):601–3.
- 26. Overstreet R, Jacobson L, Handley R. The effect of calcium on the absorption of potassium by barley roots. Plant Physiol. 1952;27(3):583.
- 27. Jayasinghe C, Pinnawala UC, Rathnayaka T, Waduge V. Annual committed effective dosage from natural radionuclides by ingestion of local food growing in mineral mining area, Sri Lanka. Environ Geochem Health. 2019;1–10.

Rev. Soc. cient. Parag. 2019;24(2):262-273

- 28. Akhter P, Rahman K, Orfi SD, Ahmad N. Radiological impact of dietary intakes of naturally occurring radionuclides on Pakistani adults. Food Chem Toxicol. 2007 Feb;45(2):272–7.
- 29. United Nations Scientific Committee on the Effects of Atomic Radiation UNSCEAR. Sources and effects of ionizing radiation: sources. Vol. 1. United Nations Publications; 2000.
- 30. Scheibel V. Determinação das Doses Efetivas por Ingestão de Farinhas de Cereais através da Espectrometria de Raios Gama. 2006. [Dissertação Mestrado]. Londrina: Universidade Estadual de Londrina, PR; 2006.
- 31. Amancio OMS, Hilário MOE. Deficiência de cobre. Rev nutr PUCCAMP. 1989;2(1):98–114.
- 32. Sargentelli V, Mauro AE, Massabni AC. Aspectos do metabolismo do cobre no homem. Quim Nova. 1996;19(3):290–3.
- 33. Unscear S. Effects of Ionizing Radiation. United Nations: New York, 2000;453–87.